

GEOMETRY OF FREE CYCLIC SUBMODULES OVER TERNIONS

HANS HAVLICEK, ANDRZEJ MATRAŚ, AND MARK PANKOV

Dedicated to Walter Benz on the occasion of his 80th birthday.

ABSTRACT. Given the algebra T of ternions (upper triangular 2×2 matrices) over a commutative field F we consider as set of points of a projective line over T the set of all free cyclic submodules of T^2 . This set of points can be represented as a set of planes in the projective space over F^6 . We exhibit this model, its adjacency relation, and its automorphic collineations. Despite the fact that T admits an F -linear antiautomorphism, the plane model of our projective line does not admit any duality.

1. INTRODUCTION

Note that all our rings are associative, with a unit element $1 \neq 0$, which acts unitaly on modules, and is inherited by subrings.

One of the crucial tasks in ring geometry is to find a “good” definition of the *projective line* over a ring R . In terms of left homogeneous coordinates a *point* of such a line should be a *cyclic submodule* $R(a, b)$ of the free R -left module R^2 . But which pairs (a, b) should be allowed to be generators of points? Indeed, there are different definitions and we refer to [16] and [22] for detailed discussion which includes also “higher-dimensional” spaces.

The aim of the present paper is to exhibit the interplay between two notions: On the one hand we consider as *points* the *free cyclic submodules* $R(a, b)$ of R^2 . On the other hand we have the distinguished subset of *unimodular points*, *i. e.*, points of the form $R(a, b)$ such that there are $x, y \in R$ with $ax + by = 1$. Several authors consider as points only our unimodular points. Often also extra conditions on the coordinate ring (like R being of stable rank 2) can be found; see [7], [14], or [22]. At the other extreme, in *projective lattice geometry* any cyclic (or: 1-generated) submodule $R(a, b) \subset R^2$ is called a point, whereas our points are called *free points* in this context [8, p. 1129]; see also [12].

For any ring R , the submodule $R(1, 0)$ is a unimodular point, but the existence of non-unimodular points cannot be guaranteed. An easy example is the projective line over any field. All its points are unimodular. So it seems necessary to restrict ourselves to a class of rings for which non-unimodular points do exist. One such class is formed by the algebras of ternions over commutative fields. There are several articles which describe the geometry over ternions based on unimodular points [1], [2], [3], [10], [11] in great detail, but little seems to be known about the properties of the remaining (non-unimodular) points [13], [20], [21].

Results and notions which are used without further reference can be found, for example, in [19].

2. CYCLIC SUBMODULES

Let F be a (commutative) field and T be the ring of *ternions*, *i. e.*, upper triangular 2×2 matrices

$$\begin{pmatrix} x & y \\ 0 & z \end{pmatrix},$$

over F . We shall often identify $x \in F$ with the ternion xI , where $I \in T$ denotes the 2×2 identity matrix. In terms of this identification F equals the centre of T . The ring T is a non-commutative *three-dimensional* algebra over F , a fact which is reflected in its name¹. A nice algebraic characterisation (avoiding associativity) can be found in [17]: Every non-commutative unital left-alternative F -algebra of dimension three is isomorphic to the algebra of ternions over F .

By [13, Lemma 2], the non-zero cyclic submodules of the free T -left module T^2 fall into five orbits under the action of the group $\mathrm{GL}_2(T)$. Below we give one representative for each orbit:

$$(1) \quad X_0 := T \left[\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \right] = \left\{ \left[\begin{pmatrix} x & y \\ 0 & z \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \right] \mid x, y, z \in F \right\},$$

$$(2) \quad Y_0 := T \left[\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right] = \left\{ \left[\begin{pmatrix} 0 & y \\ 0 & z \end{pmatrix}, \begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix} \right] \mid x, y, z \in F \right\},$$

$$(3) \quad \alpha_0 := T \left[\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \right] = \left\{ \left[\begin{pmatrix} 0 & y \\ 0 & z \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \right] \mid y, z \in F \right\},$$

$$(4) \quad \beta_0 := T \left[\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \right] = \left\{ \left[\begin{pmatrix} x & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \right] \mid x \in F \right\},$$

$$(5) \quad \gamma_0 := T \left[\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \right] = \left\{ \left[\begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \right] \mid x \in F \right\}.$$

An arbitrary submodule from the orbit of X_0 has the form $X = X_0S$ with $S \in \mathrm{GL}_2(T)$, and will be called an *X-submodule* for short. Similarly the other types of cyclic submodules are called *Y-submodules*, *α -submodules*, and so on. Submodules of the first two types are free, whence they are elements of the set of points. The unimodular points are the *X*-submodules, the non-unimodular points are the *Y*-submodules of T^2 . The remaining three types are torsion.

Any 2×2 matrix S over T can be considered as a 4×4 matrix over F of the block form

$$(6) \quad \left(\begin{array}{cc|cc} a_{11} & a_{12} & b_{11} & b_{12} \\ 0 & a_{22} & 0 & b_{22} \\ \hline c_{11} & c_{12} & d_{11} & d_{12} \\ 0 & c_{22} & 0 & d_{22} \end{array} \right).$$

We have $S \in \mathrm{GL}_2(T)$ if, and only if, its determinant (as matrix over F) satisfies

$$(7) \quad \det S = (a_{22}d_{22} - b_{22}c_{22})(a_{11}d_{11} - b_{11}c_{11}) \neq 0,$$

because invertibility of S over F implies that S^{-1} can be partitioned into four ternions as in (6). Hence it is easy to determine whether or not a 2×2 matrix over T is invertible or not. See [4] and [11, pp. 9–10] for more results on invertibility and the actual inversion of matrices over ternions.

¹According to [3] the term “ternions” is due to E. Study (1889). It is worth noting that J. Petersen used the same phrase in a different meaning already in 1885, namely as a name for a three-dimensional commutative algebra over the real numbers [18, 3.2].

By assuming S to be invertible, we obtain from (1)–(5) and (7) the following general form for X -submodules, Y -submodules, α -submodules, and so on:

$$(8) \quad X = \left\{ \left[\begin{pmatrix} a_{11}x & a_{12}x + a_{22}y \\ 0 & a_{22}z \end{pmatrix}, \begin{pmatrix} b_{11}x & b_{12}x + b_{22}y \\ 0 & b_{22}z \end{pmatrix} \right] \mid x, y, z \in F \right\} \\ \text{with } (a_{11}, b_{11}), (a_{22}, b_{22}) \neq (0, 0),$$

$$(9) \quad Y = \left\{ \left[\begin{pmatrix} 0 & a_{22}y + c_{22}x \\ 0 & a_{22}z \end{pmatrix}, \begin{pmatrix} 0 & b_{22}y + d_{22}x \\ 0 & b_{22}z \end{pmatrix} \right] \mid x, y, z \in F \right\} \\ \text{with } (a_{22}d_{22} - b_{22}c_{22}) \neq 0,$$

$$(10) \quad \alpha = \left\{ \left[\begin{pmatrix} 0 & a_{22}y \\ 0 & a_{22}z \end{pmatrix}, \begin{pmatrix} 0 & b_{22}y \\ 0 & b_{22}z \end{pmatrix} \right] \mid y, z \in F \right\} \text{ with } (a_{22}, b_{22}) \neq (0, 0),$$

$$(11) \quad \beta = \left\{ \left[\begin{pmatrix} a_{11}x & a_{12}x \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} b_{11}x & b_{12}x \\ 0 & 0 \end{pmatrix} \right] \mid x \in F \right\} \text{ with } (a_{11}, b_{11}) \neq (0, 0),$$

$$(12) \quad \gamma = \left\{ \left[\begin{pmatrix} 0 & a_{22}x \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & b_{22}x \\ 0 & 0 \end{pmatrix} \right] \mid x \in F \right\} \text{ with } (a_{22}, b_{22}) \neq (0, 0).$$

Conversely, any subset of T^2 as in (8)–(12) is easily seen to be a submodule of the appropriate type.

3. REPRESENTATION OF POINTS

The unimodular points of T^2 can be represented as subspaces of an F -vector space as follows. Let U be a *faithful* right module over T of F -dimension r . (Recall that $F \subset T$ due to our identification.) It will be convenient to write $xu := ux$ for all $u \in U$ and all $x \in F$. So elements of F may act on U from either side, whereas proper ternions act from the right hand side only. Then $U \times U$ is a right T -module in the usual way and at the same time an F -vector space of dimension $2r$. The assignment

$$(13) \quad T(A, B) \mapsto \{(uA, uB) \mid u \in U\}$$

defines an *injective* map Φ_U , say, of the set of unimodular points into the Grassmannian $\mathcal{G}_r(U \times U)$ of r -dimensional subspaces of $U \times U$. This is a direct consequence of more general results from [5, Theorem 4.2] and [14, pp. 805–806]. Like many authors we adopt the projective point of view: The elements of the Grassmannian $\mathcal{G}_r(U \times U)$ will be identified with the corresponding $(r - 1)$ -flats (projective subspaces) of the $(2r - 1)$ -dimensional projective space on $U \times U$ (over F).

The easiest example is obtained by choosing $U := F^2$ and by defining the right action of a ternion on a row $(u_1, u_2) \in F^2$ as the usual matrix multiplication. It is convenient to identify here the pair $((u_1, u_2), (v_1, v_2)) \in U \times U$ with the row vector $(u_1, u_2, v_1, v_2) \in F^4$. This leads to a well known model for the set of unimodular points as a set of lines (1-flats) in the three-dimensional projective space over F : The images of the unimodular points under Φ_{F^2} are precisely those lines of this projective space which meet the line generated by $(0, 1, 0, 0)$ and $(0, 0, 0, 1)$ at an arbitrary point (1-dimensional subspace, 0-flat). In other words (cf., *e. g.*, the table in [15, p. 30]), the unimodular points are represented by the lines of a *special linear complex* without its axis. A proof can be found in [5, Example 5.5] or [7, p. 239] (up to a permutation of coordinates due to the usage of lower triangular matrices). For the reader's convenience let us sketch the easy proof. The point X from (8) is mapped under Φ_{F^2} to the subspace comprising all vectors of the form

$$\left[(u_1, u_2) \begin{pmatrix} a_{11}x & a_{12}x + a_{22}y \\ 0 & a_{22}z \end{pmatrix}, (u_1, u_2) \begin{pmatrix} b_{11}x & b_{12}x + b_{22}y \\ 0 & b_{22}z \end{pmatrix} \right]$$

with variable $u_1, u_2, x, y, z \in F$. This subspace is spanned by the linearly independent vectors

$$(a_{11}, a_{12}, b_{11}, b_{12}) \quad \text{and} \quad (0, a_{22}, 0, b_{22}),$$

whence it is a line. The same representation was obtained in several papers on projective geometry over ternions [2, p. 157], [10, pp. 128–132], [11, Part C]. The mapping Φ_{F^2} can be extended to a mapping of non-unimodular points by following (13). Indeed, the Y -submodule (9) is mapped to the subspace

$$\left[(u_1, u_2) \begin{pmatrix} 0 & a_{22}y + c_{22}x \\ 0 & a_{22}z \end{pmatrix}, (u_1, u_2) \begin{pmatrix} 0 & b_{22}y + d_{22}x \\ 0 & b_{22}z \end{pmatrix} \right]$$

with variable $u_1, u_2, x, y, z \in F$. But this subspace does not depend on the choice of Y , because it is spanned by the vectors $(0, 1, 0, 0)$ and $(0, 0, 1, 0)$. Thus, in projective terms, *all* non-unimodular points are mapped to one line, namely the axis of the special linear complex we encountered before. Hence this extended map is no longer injective and therefore of little use.

In order to obtain an injective representation of unimodular and non-unimodular points, we make use of the *right regular representation* of T , that is we let $U = T$. As the mapping Φ_T is the identity, there will be no need to write it down explicitly below. There is the natural identification Φ of the free left module T^2 with the vector space F^6 :

$$(14) \quad \left[\begin{pmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{pmatrix}, \begin{pmatrix} b_{11} & b_{12} \\ 0 & b_{22} \end{pmatrix} \right] \xrightarrow{\Phi} (a_{11}, b_{11}, a_{22}, b_{22}, a_{12}, b_{12}).$$

We put x_1, x_2, \dots, x_6 for the coordinates in F^6 . Any 2×2 matrix S over T defines a linear mapping $g : T^2 \rightarrow T^2$ and a linear mapping $f : F^6 \rightarrow F^6$ which is characterised by $\Phi \circ g = f \circ \Phi$. If S is given as in (6) the associated matrix of f has the block form

$$(15) \quad \left(\begin{array}{cc|cc|cc} a_{11} & b_{11} & 0 & 0 & a_{12} & b_{12} \\ c_{11} & d_{11} & 0 & 0 & c_{12} & d_{12} \\ \hline 0 & 0 & a_{22} & b_{22} & 0 & 0 \\ 0 & 0 & c_{22} & d_{22} & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & a_{22} & b_{22} \\ 0 & 0 & 0 & 0 & c_{22} & d_{22} \end{array} \right).$$

Equation (7) shows that the matrix in (15) is invertible if, and only if, the matrix S is invertible.

Denote by $\mathcal{G}_k(F^6) =: \mathcal{G}_k$ the Grassmannian consisting of k -dimensional subspaces of F^6 , $k \in \{1, 2, 3, 4, 5\}$. Like before the elements of this Grassmannian will be identified with the corresponding $(k-1)$ -flats of the 5-dimensional projective space over F (points, lines, planes, solids, hyperplanes). We write \mathcal{G} for the set formed by the Φ -images of all non-zero cyclic submodules. For each $i \in \{X, Y, \alpha, \beta, \gamma\}$ we denote by \mathcal{G}_i the set of the Φ -images of all submodules of type i . So,

$$(16) \quad \mathcal{G} = \mathcal{G}_X \cup \mathcal{G}_Y \cup \mathcal{G}_\alpha \cup \mathcal{G}_\beta \cup \mathcal{G}_\gamma.$$

Since every element $S \in \text{GL}_2(T)$ induces a linear automorphism of F^6 according to (15), the Φ -images of submodules of the same type have the same dimension. From (1)–(5), we obtain

$$\mathcal{G}_X \cup \mathcal{G}_Y \subset \mathcal{G}_3, \quad \mathcal{G}_\alpha \subset \mathcal{G}_2, \quad \mathcal{G}_\beta \cup \mathcal{G}_\gamma \subset \mathcal{G}_1.$$

Hence all non-zero cyclic submodules of T^2 are represented by non-zero subspaces of F^6 . Since $\Phi : T^2 \rightarrow F^6$ is injective, the following holds trivially:

Proposition 1. *The Φ -images of distinct cyclic submodules of T^2 are distinct subspaces of F^6 .*

In particular, the Φ -images of distinct free cyclic submodules are distinct planes of F^6 .

4. STRUCTURE OF \mathcal{G}

Even though we aim at describing $\mathcal{G}_X \cup \mathcal{G}_Y$, *i. e.*, the Φ -images of free cyclic submodules of T^2 , we shall exhibit the entire set \mathcal{G} from (16), because the remaining elements of \mathcal{G} will turn out useful. First we recall some basic notions for Grassmannians \mathcal{G}_k , $k \in \{1, 2, 3, 4, 5\}$. If V and W are subspaces of F^6 with $V \subset W$ then $[V, W]_k$ denotes the subset of \mathcal{G}_k formed by all k -dimensional subspaces which contain V and are contained in W . If moreover $\dim V = k - 1$ and $\dim W = k + 1$ then $[V, W]_k$ is called a *pencil* of \mathcal{G}_k .

In our further investigation the solids

$$(17) \quad J \text{ defined by the conditions } x_3 = x_4 = 0,$$

$$(18) \quad K \text{ defined by the conditions } x_1 = x_2 = 0,$$

and their intersection, namely the line

$$(19) \quad L \text{ defined by the conditions } x_1 = x_2 = x_3 = x_4 = 0$$

will play a crucial role. Furthermore, in the solid K we have the hyperbolic quadric

$$(20) \quad H \text{ defined by the conditions } x_1 = x_2 = \det((x_3, x_4), (x_5, x_6)) = 0.$$

J and K (and henceforth L) are invariant subspaces of any linear bijection of F^6 which arises from $S \in \text{GL}_2(T)$ according to (15). This can be read off immediately from the rows of the matrix in (15). Furthermore, also the hyperbolic quadric H is easily seen to be invariant under any such linear bijection.

Proposition 2. *The following assertions are fulfilled:*

- (1) \mathcal{G}_γ coincides with the set of all points of the line L . The line L is a generator of the hyperbolic quadric H .
- (2) \mathcal{G}_β coincides with the set of all points of the solid J which are off the line L .
- (3) \mathcal{G}_α is that regulus of the hyperbolic quadric H which does not contain L .

Proof. The first two assertions hold because of (12) and (11). To show the last assertion we first notice that $\Phi(\gamma_0) = \Phi(\alpha_0) \cap L$ and that $\Phi(\alpha_0)$ is a line of the hyperbolic quadric H . As S varies in $\text{GL}_2(T)$ the point $\Phi(\gamma_0 S) = \Phi(\alpha_0 S) \cap L$ ranges in \mathcal{G}_γ , whence the line $\Phi(\alpha_0 S)$ ranges in that regulus on H which does not contain L . \square

Proposition 3. *\mathcal{G}_Y is a pencil of planes, namely $[L, K]_3$.*

Proof. Let Y be a free submodule as in (9). Hence $\Phi(Y)$ is contained in the solid K defined in (18). Letting $z = 0$ in (9) shows that the line L is contained in the plane $\Phi(Y)$. So we have $\mathcal{G}_Y \subset [L, K]_3$.

Each plane $N \subset K$ satisfies conditions

$$x_1 = x_2 = 0, \quad ax_3 + bx_4 + cx_5 + dx_6 = 0$$

for some $a, b, c, d \in F$ not all 0. If this plane contains L then $cx_5 + dx_6 = 0$ for all $x_5, x_6 \in F$. Hence $c = d = 0$ and $(a, b) \neq (0, 0)$. We define $a_{22} := -b$ and $b_{22} := a$. So there exist $c_{22}, d_{22} \in F$ such that $(a_{22}d_{22} - b_{22}c_{22}) \neq 0$. By the remark after (8)–(12), there exists a free submodule Y with $\Phi(Y) = N$. Therefore, \mathcal{G}_Y coincides with the pencil $[L, K]_3$. \square

Proposition 4. \mathcal{G}_X consist of all planes M of the Grassmannian \mathcal{G}_3 which satisfy the following two conditions:

$$(21) \quad M \cap J \text{ is a line other than } L.$$

$$(22) \quad M \cap K \text{ is a line belonging to the regulus } \mathcal{G}_\alpha.$$

Proof. Let X be a free submodule as in (8). Letting $z = 0$ in (8) shows that $\Phi(X) \cap J$ is a line of J other than L . Letting $x = 0$ in (8) shows that $\Phi(X) \cap K$ is a line belonging to the regulus \mathcal{G}_α .

Conversely, suppose that M is a plane satisfying (21) and (22). There exists a torsion submodule β such that $\Phi(\beta)$ is a point on the line $M \cap L$, and a torsion submodule α with $\Phi(\alpha) = M \cap K$. These submodules β and α can be written as in (11) and (10) which gives coefficients $a_{11}, a_{12}, a_{22}, b_{11}, b_{12}, b_{22} \in F$ subject to conditions stated there. We use these coefficients to define a submodule X according to (8). Then $\Phi(X) = M$. \square

By the above proposition, \mathcal{G}_X is formed by all planes spanned by pairs of lines, where one of them is from J and distinct from L , and the other is from the regulus \mathcal{G}_α .

Recall that flats P, Q are said to be *incident*, in symbols PIQ , if $P \subset Q$ or $Q \subset P$. The proof of Proposition 5 below provides the lengthy description of all incident pairs (P_0, Q) for a fixed P_0 and a variable Q .

Proposition 5. *The following table displays the number of incident pairs (P_0, Q) , where $P_0 \in \mathcal{G}$ is fixed and $Q \in \mathcal{G}$ is variable:*

I	$Q \in \mathcal{G}_X$	$Q \in \mathcal{G}_Y$	$Q \in \mathcal{G}_\alpha$	$Q \in \mathcal{G}_\beta$	$Q \in \mathcal{G}_\gamma$
$P_0 \in \mathcal{G}_X$	1	0	1	$ F $	1
$P_0 \in \mathcal{G}_Y$	0	1	1	0	$ F + 1$
$P_0 \in \mathcal{G}_\alpha$	$ F ^2 + F $	1	1	0	1
$P_0 \in \mathcal{G}_\beta$	$ F + 1$	0	0	1	0
$P_0 \in \mathcal{G}_\gamma$	$ F ^2 + F $	$ F + 1$	1	0	1

Proof. We sketch the proof by completely describing all possibilities.

(a) Let P_0 in \mathcal{G}_X . Then $P_0IQ \in \mathcal{G}_X \cup \mathcal{G}_Y$ is equivalent to $Q = P_0$. $P_0IQ \in \mathcal{G}_\alpha$ is equivalent to $Q = P_0 \cap K$. $P_0IQ \in \mathcal{G}_\beta$ holds if, and only if, Q is one of the $|F|$ points on $P_0 \cap J$ other than $P_0 \cap L$. $P_0IQ \in \mathcal{G}_\gamma$ is equivalent to $Q = P_0 \cap L$.

(b) Let P_0 in \mathcal{G}_Y . Then $P_0IQ \in \mathcal{G}_X \cup \mathcal{G}_Y$ is equivalent to $Q = P_0$. $P_0IQ \in \mathcal{G}_\alpha$ holds if, and only if, Q is the only generator other than L of the hyperbolic quadric H which belongs to the plane P_0 . $P_0IQ \in \mathcal{G}_\beta$ is impossible. $P_0IQ \in \mathcal{G}_\gamma$ is equivalent to Q being one of the $|F| + 1$ points on L .

(c) Let P_0 in \mathcal{G}_α . Then $P_0IQ \in \mathcal{G}_X$ is equivalent to Q being one of the $|F|^2 + |F|$ planes of $[P_0, P_0 + J]_3$ other than $P_0 + L$. $P_0IQ \in \mathcal{G}_Y$ is equivalent to $Q = P_0 + L$. $P_0IQ \in \mathcal{G}_\alpha$ is equivalent to $Q = P_0$. $P_0IQ \in \mathcal{G}_\beta$ is impossible. $P_0IQ \in \mathcal{G}_\gamma$ is equivalent to $Q = P_0 \cap L$.

(d) Let P_0 in \mathcal{G}_β . Then $P_0IQ \in \mathcal{G}_X$ is equivalent to Q being one of the $|F| + 1$ planes of the form $P_0 + R$, where R ranges in the regulus \mathcal{G}_α . $P_0IQ \in \mathcal{G} \setminus \mathcal{G}_X$ is equivalent to $Q = P_0$.

(e) Let P_0 in \mathcal{G}_γ . Then $P_0IQ \in \mathcal{G}_\alpha$ is equivalent to $Q = R_0$, where R_0 denotes the only generator other than L of the hyperbolic quadric H through the point P_0 . (This line R_0 is also used in the next two subcases.) $P_0IQ \in \mathcal{G}_X$ is equivalent to Q being one of the $|F|^2 + |F|$ planes of $[R_0, J + R_0]_3$ other than $R_0 + L$. $P_0IQ \in \mathcal{G}_Y$ is equivalent to $Q \in \mathcal{G}_Y$, which has $|F| + 1$ elements. $P_0IQ \in \mathcal{G}_\beta$ is impossible. $P_0IQ \in \mathcal{G}_\gamma$ is equivalent to $Q = P_0$. \square

Remark 1. If we associate with each $M \in \mathcal{G}_X$ the line $M \cap J$ then a line model for the set of unimodular points is obtained. This is just the special linear complex mentioned at the beginning of Section 3 with L being the axis of the complex.

5. ADJACENCY

For any $k \in \{1, 2, 3, 4, 5\}$ the flats $Z_1, Z_2 \in \mathcal{G}_k$ are called *adjacent*, in symbols $Z_1 \sim Z_2$, if their intersection is $(k - 1)$ -dimensional. It will also be convenient to write $Z_1 \cong Z_2$ for elements which are adjacent or identical. The cases $k = 1$ and $k = 5$ are trivial, because any two distinct points and any two distinct hyperplanes are adjacent, whereas in the remaining cases the entire geometry of the Grassmannian \mathcal{G}_k can be based solely on adjacency due to the famous theorem of Chow. See, *e. g.*, Chapter 3 in [19] for more information.

Our aim is to exhibit the restriction of the adjacency relation to the sets \mathcal{G}_X , \mathcal{G}_Y , and $\mathcal{G}_X \cup \mathcal{G}_Y$. They represent the sets of unimodular points, non-unimodular points, and all points of the projective line over T . As before, the flats J, K, L , and the hyperbolic quadric H defined in (17)–(20) will be of great importance. Given $M_1, M_2 \in \mathcal{G}_X$ we have

$$(23) \quad M_1 \cong M_2 \Leftrightarrow M_1 \cap K = M_2 \cap K,$$

since for $M_1 \neq M_2$ either side of (23) is equivalent to $M_1 + M_2$ being a solid, whereas for $M_1 = M_2$ (23) holds trivially. We infer from (23) that \cong is an equivalence relation on \mathcal{G}_X . The equivalence classes are of the form

$$(24) \quad [P, P + J]_3 \setminus \{P + L\} \text{ with } P \in \mathcal{G}_\alpha.$$

So there is a one-one correspondence between these equivalence classes and the lines of the regulus \mathcal{G}_α .

We now consider the adjacency relation on $\mathcal{G}_X \cup \mathcal{G}_Y$. Each plane $M \in \mathcal{G}_X$ is adjacent with precisely one plane of \mathcal{G}_Y , namely

$$N := (M \cap K) + L.$$

As \mathcal{G}_Y is a pencil of planes, we have $N_1 \cong N_2$ for all $N_1, N_2 \in \mathcal{G}_Y$. Consequently, the pencil $[L, K]_3$ and the subsets

$$(25) \quad [P, P + J]_3 \text{ with } P \in \mathcal{G}_\alpha$$

are the *cliques* of $\mathcal{G}_X \cup \mathcal{G}_Y$ with respect to adjacency. That means, these are the maximal subsets of $\mathcal{G}_X \cup \mathcal{G}_Y$ for which any two distinct elements are adjacent. It is well known that the elements of any clique (25) and the pencils contained in it can be considered as the “points” and “lines” of a projective plane. Thus any equivalence class from (24) can be regarded as a *punctured projective plane*, *i. e.*, a projective plane with one point removed. Going over from \mathcal{G}_X to $\mathcal{G}_X \cup \mathcal{G}_Y$ provides the “closure” of all these punctured projective planes.

Another advantage of $\mathcal{G}_X \cup \mathcal{G}_Y$ over \mathcal{G}_X is *connectedness* with respect to \sim , *i. e.*, given any $Z, Z' \in \mathcal{G}_X \cup \mathcal{G}_Y$ there exists a finite sequence Z_0, Z_1, \dots, Z_r of planes from $\mathcal{G}_X \cup \mathcal{G}_Y$ such that

$$Z = Z_0 \sim Z_1 \sim \dots \sim Z_r = Z'.$$

The minimal $r \geq 0$ for which such a sequence exists is said to be the *distance* between Z and Z' . Any two *distinct* planes $M_1, M_2 \in \mathcal{G}_X$ are either at distance 1 (adjacent) or at distance 3, because for $M_1 \not\sim M_2$ we have

$$M_1 \sim (M_1 \cap K) + L \sim (M_2 \cap K) + L \sim M_2,$$

and this is the only shortest sequence from M_1 to M_2 . Yet, this distance function is of restricted use, since in the latter case $M_1 \cap M_2$ may be 0 (the “empty flat”) or a single point.

Any bijection of $\mathcal{G}_X \cup \mathcal{G}_Y$ onto itself which preserves adjacency in both directions will be called an *adjacency preserver* of $\mathcal{G}_X \cup \mathcal{G}_Y$. The straightforward proof of the following result is left to the reader:

Proposition 6. *Let $\mu : \mathcal{G}_\alpha \rightarrow \mathcal{G}_\alpha$ be a bijection. For each $P \in \mathcal{G}_\alpha$ we choose a bijection*

$$(26) \quad \psi_P : [P, P + J]_3 \rightarrow [\mu(P), \mu(P) + J]_3 \quad \text{such that} \quad P + L \mapsto \mu(P) + L.$$

Then the mapping

$$(27) \quad \lambda : \mathcal{G}_X \cup \mathcal{G}_Y \rightarrow \mathcal{G}_X \cup \mathcal{G}_Y : Z \mapsto \psi_P(Z) \quad \text{if} \quad Z \in [P, P + J]_3$$

is a well-defined adjacency preserver of $\mathcal{G}_X \cup \mathcal{G}_Y$. Conversely, every adjacency preserver of $\mathcal{G}_X \cup \mathcal{G}_Y$ arises in this way.

Due to the condition on ψ_P stated at the end of formula (26) we obtain the following:

Corollary 1. *Every adjacency preserver λ of $\mathcal{G}_X \cup \mathcal{G}_Y$ satisfies $\lambda(\mathcal{G}_X) = \mathcal{G}_X$ and $\lambda(\mathcal{G}_Y) = \mathcal{G}_Y$.*

If we impose the much stronger requirement that a bijection of $\mathcal{G}_X \cup \mathcal{G}_Y$ onto itself should map pencils of planes onto pencils of planes then all bijections from (26) have to be collineations between the underlying projective planes, but there need not be any relationship between these collineations. So even here we obtain transformations as in (27) that do not really deserve our interest.

Still, we have the following description of adjacency preservers which arise from semilinear bijections. Recall that via the bijection (14) the points of the projective line over T correspond to the planes belonging to $\mathcal{G}_X \cup \mathcal{G}_Y$.

Theorem 1. *For all mappings $g : T^2 \rightarrow T^2$ and all mappings $f : F^6 \rightarrow F^6$ such that $\Phi \circ g = f \circ \Phi$ the following statements are equivalent.*

- (i) *g is a T -semilinear bijection of the module T^2 .*
- (ii) *f is an F -semilinear bijection of the vector space F^6 satisfying $f(\mathcal{G}_X \cup \mathcal{G}_Y) = \mathcal{G}_X \cup \mathcal{G}_Y$.*
- (iii) *f is an F -semilinear bijection of the vector space F^6 satisfying $f(\mathcal{G}_X) = \mathcal{G}_X$.*
- (iv) *f is an F -semilinear bijection of the vector space F^6 stabilising the solid J and the hyperbolic quadric H which are defined by $x_3 = x_4 = 0$ and $x_1 = x_2 = \det((x_3, x_4), (x_5, x_6))$, respectively.*

We postpone the proof until we have established two lemmas.

Lemma 1. *A line Q has the property that for all $M \in \mathcal{G}_X$ the intersection $Q \cap M$ is a point if, and only if, Q belongs to the opposite regulus of \mathcal{G}_α .*

Proof. (a) Let Q be a line such that $Q \cap M$ is a point for all $M \in \mathcal{G}_X$. We first choose a fixed $P \in \mathcal{G}_\alpha$ and define the hyperplane $W := P + J$. Then all but one plane of $[P, W]_3$ belong to \mathcal{G}_X ; the only exception is $P + L \in \mathcal{G}_Y$. We claim that $Q \cap P$ contains a point. Assume to the contrary that this were not the case. So, for all $M \in [P, W]_3 \cap \mathcal{G}_X$, the point $Q \cap M$ would be off the line P . This would imply that all such M could be written as $M = P + (M \cap Q)$, whence all of them would be contained in the solid $P + Q$, a contradiction.

Now, as P ranges in the regulus \mathcal{G}_α , we see that Q is a line of the regulus opposite to \mathcal{G}_α .

(b) The converse is obviously true. □

Lemma 2. *A solid V has the property that for all $M \in \mathcal{G}_X$ the intersection $V \cap M$ is a line if, and only if, either $V = J$ or $V = K$.*

Proof. (a) Let V be a solid such that $V \cap M$ is a line for all $M \in \mathcal{G}_X$. We first choose a fixed $P \in \mathcal{G}_\alpha$ and define the hyperplane $W := P + J$. Then all but one planes of $[P, W]_3$ belong to \mathcal{G}_X ; the only exception is $P + L \in \mathcal{G}_Y$. We claim that

$$(28) \quad V \subset W \Leftrightarrow P \not\subset V.$$

On the one hand, for $V \subset W$ we obtain $P \not\subset V$, since otherwise there would exist an $M \in [P, V]_3 \cap \mathcal{G}_X$, and $M \cap V = M$ would not be a line. On the other hand, for $V \not\subset W$ the intersection $V \cap W$ is a plane. We have $P \subset V$, since otherwise there would exist an $M \in [P, W]_3 \cap \mathcal{G}_X$ with $M \cap (V \cap W) = M \cap V$ being a point. For the rest of the proof we distinguish two cases:

Case 1: There exists a line of \mathcal{G}_α , say P_1 , with $P_1 \subset V$. There exist two further lines $P_2, P_3 \in \mathcal{G}_\alpha$ and three mutually skew planes $M_1, M_2, M_3 \in \mathcal{G}_X$ with $M_i \cap K = P_i$ for $i \in \{1, 2, 3\}$. Through each point of the plane M_1 there is a unique line which meets each of the planes M_2 and M_3 at a point. Within the solid V we have a similar result about the mutually skew lines $M_1 \cap V, M_2 \cap V, M_3 \cap V$: Through each point of the line $M_1 \cap V = P_1$ there is a unique line which meets each of the lines $M_2 \cap V$ and $M_3 \cap V$ at a point. Furthermore, through each point of P_1 there is a unique line of the regulus which is opposite to \mathcal{G}_α , and this line meets each of the lines $P_2 \subset M_2$ and $P_3 \subset M_3$ at a point. We combine these three observations and infer that all lines of the opposite regulus of \mathcal{G}_α are contained in V , that $P \subset V$ for all $P \in \mathcal{G}_\alpha$, and finally that $V = K$.

Case 2: For all $P \in \mathcal{G}_\alpha$ holds $P \not\subset V$. From (28) (with W to be replaced by $P + J$) we infer $V \subset P + J$ for all $P \in \mathcal{G}_X$. Due to $\bigcap_{P \in \mathcal{G}_X} (P + J) = J$, now there holds $V = J$. \square

Proof of Theorem 1. (i) \Rightarrow (ii): The accompanying automorphism of g is, like any automorphism of T , the product of an automorphism σ of F (acting entrywise on T) followed by an inner automorphism of T [9, Theorem 6.6]. By virtue of this result, a straightforward calculation shows that $f = \Phi \circ g \circ \Phi^{-1}$ is an F -semilinear bijection with respect to σ . The assertion $f(\mathcal{G}_X) = \mathcal{G}_X$ is obviously true.

(ii) \Rightarrow (iii): The F -semilinear bijection f is an adjacency preserver on $\mathcal{G}_X \cup \mathcal{G}_Y$. By Corollary 1 we have $f(\mathcal{G}_X) = \mathcal{G}_X$.

(iii) \Rightarrow (iv): We infer from Lemma 1 that the regulus opposite to \mathcal{G}_α is fixed, as a set of lines, under f . Therefore also the hyperbolic quadric H and the solid K are invariant under f . From $f(K) = K$ and Lemma 2 follows $f(J) = J$.

(iv) \Rightarrow (i): The F -semilinear bijection f can be written as a product of three F -semilinear bijections f_1, f_2, f_3 as follows:

The first bijection is $f_1 : (x_1, x_2, \dots, x_6) \mapsto (\sigma(x_1), \sigma(x_2), \dots, \sigma(x_6))$, where σ is the automorphism of F accompanying f . Then $g_1 := \Phi^{-1} \circ f_1 \circ \Phi$ is clearly a T -semilinear bijection of T^2 .

The second mapping is the unique linear bijection $f_2 : F^6 \rightarrow F^6$ which fixes all vectors of J and permutes the lines of the regulus opposite to \mathcal{G}_α in the same way as $f \circ f_1^{-1}$. All lines of \mathcal{G}_α are two-dimensional f_2 -invariant subspaces. The matrix of f_2 can therefore be written in block form as

$$\begin{pmatrix} I & 0 & 0 \\ 0 & bI & aI \\ 0 & 0 & I \end{pmatrix} \in \text{GL}_6(F),$$

where I denotes the 2×2 identity matrix. The homothety

$$g_2 : T^2 \rightarrow T^2 : (A, B) \mapsto \begin{pmatrix} 1 & a \\ 0 & b \end{pmatrix} (A, B)$$

is T -semilinear (with respect to an inner automorphism) and an easy calculation shows $\Phi \circ g_2 = f_2 \circ \Phi$.

The third mapping is the linear bijection $f_3 := f \circ f_1^{-1} \circ f_2^{-1}$. It has J, K and all lines of the regulus opposite to \mathcal{G}_α as invariant subspaces. This implies that the matrix of f_3 can be written as in (15), whence the invertible matrix (6) defines $g_3 : T^2 \rightarrow T^2$ with the property $\Phi \circ g_3 = f_3 \circ \Phi$.

Finally, from $f = f_3 \circ f_2 \circ f_1$ follows that $g = g_3 \circ g_2 \circ g_1$ is a T -semilinear bijection. \square

The previous theorem describes all automorphic collineations of $\mathcal{G}_X \cup \mathcal{G}_Y$ and also of \mathcal{G}_X . It is in the spirit of results from [23], [24], and [25].

Remark 2. A duality of the projective space on F^6 maps any Grassmannian \mathcal{G}_k , $k \in \{1, 2, 3, 4, 5\}$, bijectively onto the Grassmannian \mathcal{G}_{6-k} , and it preserves adjacency in both directions. In particular, \mathcal{G}_3 is mapped onto itself. However, the following holds: *Neither $\mathcal{G}_X \cup \mathcal{G}_Y$ nor \mathcal{G}_X is fixed, as a set of planes, under a duality of the projective space on F^6 .*

Let δ be a duality fixing $\mathcal{G}_X \cup \mathcal{G}_Y$. Hence the restriction of δ to $\mathcal{G}_X \cup \mathcal{G}_Y$ is an adjacency preserver. By Corollary 1, we obtain $\delta(\mathcal{G}_X) = \mathcal{G}_X$.

Assume to the contrary that there exists a duality fixing \mathcal{G}_X . This duality maps the lines Q , described in Lemma 1, to the solids V , described in Lemma 2, in a bijective way. We obtain a contradiction, since there are $|F| + 1 > 2$ such lines, but only two such solids.

Remark 3. Every *Jordan automorphism* of an arbitrary ring R of stable rank 2 defines a bijective mapping of the set of unimodular points of R^2 onto itself. See [7, 4.2] or [14, p. 832] for further details and, in particular, the rather complicated definition of such a mapping. For our F -algebra of ternions the situation is less intricate, since every Jordan automorphism of T is either an automorphism or an antiautomorphism; moreover, it is F -semilinear [9, Theorem 6.6]. We already referred to this result in the proof of Theorem 1, where we described all automorphisms of T . We add, for the sake of completeness, that an antiautomorphism of T is given by

$$\begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \mapsto \begin{pmatrix} z & y \\ 0 & x \end{pmatrix}.$$

The mappings of unimodular points arising from antiautomorphisms appear also in [3] for the ternions over the real numbers and in [6] in a more general setting.

In terms of the line model from the beginning of Section 3 any mapping ω on unimodular points arising from an antiautomorphism of T is induced by a duality which preserves the axis of the special linear complex. All lines through one of the points of this axis go over to all lines in one of the planes through the axis. From Remark 1 and Proposition 4, in our model we obtain from ω a bijection ξ of \mathcal{G}_X onto itself which *does not preserve adjacency*. Indeed, there are $M_1, M_2 \in \mathcal{G}_X$ with $M_1 \cap M_2 \in \mathcal{G}_\alpha$ for which the lines $M_1 \cap K$, $M_2 \cap K$, and L are not coplanar. Therefore $\xi(M_1) \cap \xi(M_2) \in \mathcal{G}_\beta$ is only a single point. However, ξ preserves (unordered) pairs of skew planes, since these correspond via Φ to *distant* unimodular points (in the terminology of [7] and [14]) and ω preserves (unordered) pairs of distant points.

Also, there does not seem to be a natural extension of ω to non-unimodular points. This is in sharp contrast to the observation in the introduction of [12] that non-unimodular points are “indispensable” for an arbitrary semilinear mapping to define a morphism of projective spaces over rings.

Acknowledgements. This work was carried out within the framework of the Scientific and Technological Cooperation Poland-Austria 2010–2011.

REFERENCES

- [1] H. Beck. Eine Cremonasche Raumgeometrie. *J. Reine Angew. Math.*, 175:129–158, 1936.

- [2] H. Beck. Über Ternionen in der Geometrie. *Math. Z.*, 40(1):509–520, 1936.
- [3] W. Benz. Über eine Cremonasche Raumgeometrie. *Math. Nachr.*, 80:225–243, 1977.
- [4] W. Benz. Zur Umkehrung von Matrizen im Bereich der Ternionen. *Mitt. Math. Ges. Hamburg*, 10(7):509–512, 1979.
- [5] A. Blunck and H. Havlicek. Projective representations I. Projective lines over rings. *Abh. Math. Sem. Univ. Hamburg*, 70:287–299, 2000.
- [6] A. Blunck and H. Havlicek. The dual of a chain geometry. *J. Geom.*, 72:27–36, 2001.
- [7] A. Blunck and A. Herzer. *Kettengeometrien – Eine Einführung*. Shaker Verlag, Aachen, 2005.
- [8] U. Brehm, M. Greferath, and S. E. Schmidt. Projective geometry on modular lattices. In F. Buekenhout, editor, *Handbook of Incidence Geometry*. Elsevier, Amsterdam, 1995.
- [9] W. L. Chooi and M. H. Lim. Coherence invariant mappings on block triangular matrix spaces. *Linear Algebra Appl.*, 346:199–238, 2002.
- [10] J. Depunt. Sur la géométrie ternionienne dans le plan. *Bull. Soc. Math. Belg.*, 11:123–133, 1959.
- [11] J. Depunt. Grondslagen van de analytische projectieve ternionenmeetkunde van het platte vlak. *Verh. Konink. Acad. Wetensch. Lett. Schone Kunst. België, Kl. Wetensch.*, 22(63):99 pp., 1960.
- [12] C.-A. Faure. Morphisms of projective spaces over rings. *Adv. Geom.*, 4(1):19–31, 2004.
- [13] H. Havlicek and M. Saniga. Vectors, cyclic submodules, and projective spaces linked with ternions. *J. Geom.*, 92(1-2):79–90, 2009.
- [14] A. Herzer. Chain geometries. In F. Buekenhout, editor, *Handbook of Incidence Geometry*, pages 781–842. Elsevier, Amsterdam, 1995.
- [15] J. W. P. Hirschfeld. *Finite Projective Spaces of Three Dimensions*. Oxford University Press, Oxford, 1985.
- [16] A. Lashkhi. Harmonic maps over rings. *Georgian Math. J.*, 4:41–64, 1997.
- [17] W. Lex, V. Poneleit, and H. J. Weinert. Über die Einzigkeit der Ternionenalgebra und linksalternative Algebren kleinen Ranges. *Acta Math. Acad. Sci. Hungar.*, 35(1-2):129–138, 1980.
- [18] J. Lützen. Julius Petersen, Karl Weierstrass, Hermann Amandus Schwarz and Richard Dedekind on hypercomplex numbers. *Mat. Medd. Danske Vid. Selsk.*, 46(2):223–254, 2001. Around Caspar Wessel and the geometric representation of complex numbers (Copenhagen, 1998).
- [19] M. Pankov. *Grassmannians of Classical Buildings*, volume 2 of *Algebra and Discrete Mathematics*. World Scientific, Singapore, 2010.
- [20] M. Saniga, H. Havlicek, M. Planat, and P. Pracna. Twin “Fano-snowflakes” over the smallest ring of ternions. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 4:050, 7 pp. (electronic), 2008.
- [21] M. Saniga and P. Pracna. A Jacobson radical decomposition of the Fano-Snowflake configuration. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 4:072, 7 pp. (electronic), 2008.
- [22] F. D. Veldkamp. Projective ring planes and their homomorphisms. In R. Kaya, P. Plaumann, and K. Strambach, editors, *Rings and Geometry*, pages 289–350. D. Reidel, Dordrecht, 1985.
- [23] R. Westwick. Linear transformations on Grassmann spaces. *Pac. J. Math.*, 14:1123–1127, 1964.
- [24] R. Westwick. Linear transformations on Grassmann spaces. *Canad. J. Math.*, 21:414–417, 1969.
- [25] R. Westwick. Linear transformations on Grassmann spaces. III. *Linear Multilinear Algebra*, 2:257–268, 1974/75.

Hans Havlicek
 Institut für Diskrete Mathematik und Geometrie
 Technische Universität
 Wiedner Hauptstraße 8–10/104
 A-1040 Wien
 Austria
 havlicek@geometrie.tuwien.ac.at

Andrzej Matraś and Mark Pankov
 Faculty of Mathematics and Computer Science
 University of Warmia and Mazury

Żołnierska 14A
PL-10-561 Olsztyn
Poland
matras@uwm.edu.pl
markpankov@gmail.com